INDEX

Ainslie, G. G., 115–118.
Alfalfa weevil, 302; Sporotrichum on, 493–499.
  caterpillar, 570.
Allen, H. W., 233–235.
Allograpta obliqua, 456.
Amblymerus sp., 465.
American Association of Economic Entomologists, Proceedings, 1–232, 253–303.
  Pacific Slope Branch, Proceedings, 453–510.
Anasa andresii, 431–434.
Ancylis angulifasciana, 80–82.
Animal fumigation, 71–78.
Anticarsia gemmatilis, 521–528.
Apanteles fulvipes, 178.
Apateticus cynicus, 51–52.
  maculiventris, 52–53.
Aphid ecology, 44–50.
Aphis maidi-radicis, 236.
  pomi, 510.
  pseudobrassicæ, 67–71.
Apiary inspection, 188–192, 196–199; investigations, 282–284.
Argentine ant, municipal control, 468–472.
Army worm parasite, 377.
Arsenate of lead, freezing, 561–566.
Arsenic in bees, 364–366.
Arsenic on fruit, 90.

Baccha lemur, 456.
Bacillus amylovorus, 362.
Back, E. A., 306–311.
Batrachedra rileyi, 295–298.
Bee diseases, 192–196.
  work, 411–413.
Beekeeping, 243.
  college, 413–417.
  investigation, 406–411.
  work in Massachusetts, 417–419.
Bees, arsenic in, 364–366.
Beneficial insects, 172–178.
Bensel, G. E., 303–306.
Bilsing, W. S., 110–113.
Binocular magnifier, 370.
Blatchley, W. S., 573.
Bridwell, J. G., 472–476.
Britton, W. E., 281–282.
Bryobia pratense, 487.
Bucculatrix thurberiella, 505–510.

Cassida pallidula, 42.
Castnia therapon, 378.
Catabomba pyrastri, 454.
Cedar rust, 204–206.
Ceratitis capitata, 306–311.
Ceromasia sphenophori, 174.
Chapman, J. W., 149–167.
Cheilosia, 457.
Chermes abietis, 49.
Cherry, Syneta on, 458.
Chrysanthemum gall-fly, 461–468.
Chittenden, F. H., 571, 573.
Chrysomphalus aurantii, 489.
Cirphis unipuncta, 401.
Clover leaf tyer, 80–82.
  leaf weevil, 445.
Coccid-feeding moth, 369–370.
Cockerell, T. D. A., 235–236, 312–313.
Codling moth, 107–109; trap, 517–521.
Columbine leaf-miner, 419–424.
Comstock, J. H., 382.
Cooper, W. F., 381.
Corn-ear worm, 395–398.
Corn-root louse, 236.
Cory, E. N., 372–375, 419–424.
Cotalpa lanigera, 265.
Cotinis, 266.
Cotton, R. T., 516.
Cotton, Bucculatrix thurberiella on, 505.
Crambid notes, 115–118.
Crambus caliginosellus, 116.
  hortuellus, 116.
  luteolellus, 116.
  mutabilis, 116.
  præfectellus, 118.
  teterrellus, 116.
  trisectus, 116, 118, 119.
  vulgivagellus, 116.
Crosby, C. R., 375–376.
Cucumber beetle, 12-spotted, 366–368, 551–556.
Cucurbits, Anasa andresii on, 431–434.
Cyanide of potassium, 169–170.
Cyclocephala immaculata, 264.
Cylas formicarius, 516.

Dasyneura ulmea, 82–84.
Davidson, W. M., 454–457.
Davis, J. J., 134, 261–281.
Dean, G. A., 139–141, 245.
De Ong, E. R., 468–472.
Diabrotica 12-punctata, 366–368, 551–556.
  longicornis, 236.
Diachasma fullawayi, 307, 308, 309, 310, 311.
  tryoni, 307, 308, 309, 310, 311, 473.
Diarthronomyia hypogæa, 461–468.
Diaspis pentagona, 179.
Diatræa saccharalis, 116, 570.
  zeacolella, 116.
Diprion simile, 281–282.
Doane, R. W., 398–401.
Dove, W. E., 528–538.
Draeculacephala reticulata, 570.
Drosichh palavanica, 235.
Dusting orchard, cost, 375–376.

Dust sprays, 395–398.
Dutcher, R. A., 561–566.
Dyscinetus trachypygus, 267.

Economic entomologist, liberal training, 15–23.
Elis 5-cincta, 271.
Employment bureau, 6.
Epitrix cucumeris, 43.
fuscula, 43.
Essig, E. O., 369–370, 461–468.
Eumerus strigatus, 457.
Euphoria, 266.
Eurymerus eurytheme, 570.
Evans, A. T., 354–362.
Executive committee, 5.
Extension entomologist, work, 287–290.

Federal horticultural board, 442.
Felt, E. P., 107–109.
Fink, D. E., 366–368.
Fire blight, 362–363.
Flies, typical, 382.
Flint, W. P., 377.
Florida inspection, 224–227.
Forbes, S. A., 239–241.
Foreign pests introduced, 212–216, 216–219.
Forest insect investigations, 442.
Foulbrood, 192–196, 379.
Fracker, S. B., 253–261.
Freezing arsenate of lead, 561–566.
Fruit-fly parasites, 472.
Fruit-tree leaf Syneta, 458–461.
Fumigation, animal, 71–78.
apparatus, 222.
subterranean, 285–287.

Gargarphia solani, 42.
Gates, B. N., 417–419.
Gipsy moth conference, 315, 443.
Glaser, R. W., 149–167.
Gonepteryx rhamni, 378.
Goodwin, W. H., 91–105.
Gossard, H. A., 53–59, 59–62, 80–82, 142–144.
Gracilaria zachrysa, 213.
Graham, S. A., 549–551.
Grape-berry worm, 91–105.
Gryllotalpa gryllotalpa, 213, 219.

Harned, R. W., 295–298.
Haseman, L., 282–284, 291–294.
Hawaiian fruit-fly parasites, 472–476.
insects, 172–178.
Hayes, W. P., 23–38, 120–128.
Headlee, T. J., 84–88.
Heliophila unipuncta, 377.
Herrick, G. W., 15–23.
Hessian fly, 142–144, 236, 290, 291.
fly train, 139–141.
Holcocera iceryæella, 369–370.
Holland, E. B., 364–366.
Honey bee and blight, 59–62.
Honey bee embryology, 242.
Horticultural inspection, 200–213; law, 206–212.
Host relationships defined, 477–486.
Housefly wintering, 528–538.
Houser, J. S., 82–84, 285–287.
Howard, C. W., 236.
Howard, L. O., 172–178, 179–181, 389–392.
Hungerford, H. B., 538–549.
Hypera punctata, 445.
variabilis, 493–499.
Hyperparasitism, 480.
Hyslop, J. A., 435–438.

Imported fruit-bud weevil, 428.
poplar root weevil, 425–428.
Indian ant, introduced, 566–569.
Index committee, 12.
Insects as food, 452.
Inspection, 224–231.
facilities, 219–223.
Isosoma grande, 398, 399.
tritici, 400.
vaginicolum, 398–401.
Ixodoidea, monograph, 381.

Jalysus spinosus, 39.
Jones, T. H., 431–434.

King, G. B., 512.
King, J. L., 106.

Laboratory binocular, 370.
Lachnosterna, 261–281.
in Wisconsin, 253.
fusca, 262, 269.
grandis, 262.
larvæ as food, 389–392.
records, 253–261.
tristis, 262.
Lampa, Sven, 242.
Laphygma frugiperda, 570.
Lecanium persicæ, 369.
Leng, C. W., 573.
Lesser peach-borer, 106.
Leucotermes lucifugus, 148
Llaveia benguetensis, 235.
Ligyrus gibbosus, 264, 267.
relictus, 264, 266, 267.
Luginbill, P., 570.

Maize bill-bug, 120–128.
Magnifier, binocular, 370.
Maxson, A. C., 500–505.
Mayetiola destructor, 236.
McColloch, J. W., 23–38, 395–398, 445.
McConnell, W. R., 145–147.
McCray, A. H., 192–196, 379.
McGregor, E. A., 505–510, 556–561.
Mediterranean fruit-fly, 176–178, 306–311.
Meliana albilinea, 288.
Membership committee, 10.
Merodon equestris, 457.
Merrill, J. H., 148.
Mesogramma polita, 443.

Mesograpta geminata, 456.
marginata, 456.
Metcalf, C. L., 89.
Minnesota Reports, Index, 382.
Moore, Wm., 71–78.
Mozenette, G. F., 458–461.
Musca domestica, 147; breeding habits, 354–362; dispersion, 325–354; wintering, 528–538.

Nelson, J. A., 242.
New Jersey Mosquito Association, 314.
Nitrobenzene, 72–78, 377.
Nomenclature committee, 7.
Nominations committee, 13.
Nuttall, G. H. F., 381.

O'Byrne, F. M., 224–227.
Ohio inspection, 227–231.
O'Kane, W. C., 90.
Oncideres texana, 110–113.
Opius humilis, 307, 308, 309, 310, 311, 473.
perproximus, 473.
Osmoderma eremicola, 267.
Otiorhynchus sulcatus, 213.

Pacific Slope Branch, Proceedings, 453–510.
Paddock, F. B., 67–71.
Paragus tibialis, 457.
Parasitism, 478.
Parasitism, indirect, 480.
multiple, 485.
quaternary, 484.
secondary, 482.
tertiary, 484.
Parker, J. R., 182–187.
Parker, R. R., 147, 325–354, 438–441.
Parrott, P. J., 238.
Patch, E. M., 44–50.
Peanuts, Diabrotica injuring, 366–368.
Pearce, E. K., 382.
Pecan twig girdler, 110–113.
Pegomyia brassicæ, 136–138, 571.
cepetorum, 571.
fusciceps, 131–133, 571.
hyoscyami, 372–375.
Pellett, F. C., 317.
Pemberton, C. E., 306–311.
Pemphigus betæ, 500–505.
Pennington, W. E., 401–406.
Peridroma margaritosa, 303–306.
Periodical cicada, 53–59.
Perkinsiella saccharicida, 173–174.
Pettit, Morley, 196–199, 406–411.
Pierce, W. D., 424–431.
Phillips, E. F., 188–192, 243, 362–363, 413–417.
Phlegethontius carolina, 43.
5-maculata, 43.
Phytomyza aquilegiæ, 419–424.
Pine blister rust, 231–232.
Pipiza albipilosa, 456.
pisticoides, 456.
Pissodes strobi, 549–551.
Planidium on aphids, 510.
Poison bran mash, 245.
Polydrusus, 424–431.
impressifrons, 425–428.
sericeus, 428.
Pleurotropis epigonus, 145–147.
Polychrosis viteana, 91–105.
Polyphylla, 266.
Potassium cyanide, 169–170.
Potted plants, Sciara injuring, 538–549.
Predatism, 478.
Presidential address, 15–23; discussion, 64–67.
Privet mite, 556–561.
Prociphilus tessellata, 47.
Productive beekeeping, 317.
Prospaltella berlesei, 179–181.
Promachus fitchii, 271.
Prune, Syneta on, 458.
Pseudococcus neomexicana, var. utahensis, 313.
timberlakei, 312–313.

Quarantines, state, 299–303.
Quayle, H. J., 486–492.

Resolutions committee, 9.
Rhabdocnemis obscurus, 174–176.
Rhogas terminalis, 401–406.
Rhynchophora, 573.
Riley, W. A., 378.
Robinson, L. E., 381.
Rockwood, L. P., 493–499.
Rosewall, O. W., 443.
Rumsey, W. E., 204–206.

Saissetia oleæ, 488.
Sanders, J. G., 133, 206–212, 253–261.
Saperda tridentata, 148.
Sarcophaga aldrichii, 438–441.
Sasscer, E. R., 216–219, 219–223.
Scale insects, 382; wind dispersion, 486–492.
Schizoneura corni, 45.
Schizoneura lanigera, 47.
Schoene, W. J., 131–133, 136–138.
Sciara coprophila, 543, 547.
maggots, 538–549.
Secretary, report, 3.
Seed-corn maggot, 131–133.
Sell, R. A., 551–556.
Sesia rileyana, 40.
Siegler, E. H., 517–521.
Sladen, F. W. L., 411–413.
Small pink corn-worm, 295–298.
Smith, H. S., 477–486.
Smulyan, M. T., 510.
Solanum insects, 39–44.
Solenopsis molesta, 23–38.
Somes, M. P., 39–44.
Spaulding, Perley, 231–232.
Sphærophoria melanosa, 455.
micrura, 455.
sulphuripes, 455.

Sphenophorus callosus, 128.
maidis, 120–128.
Sporotrichum globuliferum, 493–499.
Spraying cost, 375–376, 392–395.
Stratægus antæus, 267.
Strawberry weevil, 84–88.
Sugar-cane borer, 174–176.
Sugar-cane leaf-hopper, 173–174.
Sulphur-arsenical, 84–88.
Superparasitism, 485.
Syneta albida, 458–461.
Syneta simplex, 459.
Syrphid larvæ, 89.
Syrphidæ, Economic, in California, 454–457.
Syrphus americanus, 455.
arcuatus, 455.
opinator, 455.

Talbert, T. J., 287–290.
Taylor, J. E., 299–303.
Tenuipalpus bioculatus, 556–561.
Tetrastichus giffardi, 307, 308, 309, 310, 311.
sp., 466.
Tetranychus mytilaspidis, 238.
Ticks, Monograph of, 381.
Trichius piger, 267.
Trichobaris trinotata, 43.
Triglyphothrix striatidens, 566–569.
Triphleps insidiosus, 435–438.
Turnip louse, 67–71.

Variegated cutworm, 303.
Velvet-bean caterpillar, 521–528, 570.

Warburton, C., 381.
Watson, J. R., 521–528.
Webster, F. M., 239–241.
Weevils. 573.
Weiss, H. B., 212–216, 379.
Wellhouse, Walter, 169–170.
Wenzel, O. J., 382.
Western corn root-worm, 236.
Wheat aphis, western, 182–187.
Wheat-head army worm, 288.
Wheat, Isosoma vaginicolum on, 398–401.
Wheeler, W. M., 566–569.
White grubs, 261–281; as food, 389–392, investigations, 134.
White pine weevil, 549–551.
Whitmarsh, R. D., 51–53.
Wildemuth, V. L., 570.
Wilson, R. N., 570.
Wilt, gipsy moth, 149–167, 233–235.
Wind dispersion, 486–492.
Woglum, R. S., 370, 392–395.